

# Workbook



## Table of Contents

Green's Theorem .....	2
Green's Theorem .....	2


# Green's Theorem

## Green's Theorem


### Questions:

In each of the exercises 1-3 verify Green's Theorem:  $\oint_C f dx + g dy = \iint_R (g_x - f_y) dA$ .

- 1)  $\oint_C x^2 y dx + x dy$ ; the path  $C$  is as in the illustration:



- 2)  $\oint_C (x - y^2) dx + dy$ ; the path  $C$  is as in the illustration:



- 3)  $\oint_C (x^2 - xy^3) dx + (y^2 - 2xy) dy$ ;  $C$  traces out, anticlockwise, the square with vertices:  $(0,0), (2,0), (2,2), (0,2)$ .

- 4) Compute the work done by a force field  $\mathbf{F}(x, y) = (e^x - y^3)\mathbf{i} + (\cos y + x^3)\mathbf{j}$ , on a particle which moves anticlockwise on the unit circle  $x^2 + y^2 = 1$  and completes one revolution.

- 5) Compute the integral  $\int_C \left( e^y - \tan \frac{x}{2} \right) dx + (xe^y + y \cos y^2) dy$ , where  $C$  is the clockwise union of the parts of the curves  $y = 8 - x^2$ ,  $y = x^2$  between the  $y$ -axis and their intersection in the first quadrant.

- 6) Compute the integral  $\int_C -2e^{2x-y} \cos y dx + (e^{2x-y} (\sin y + \cos y) + 2xy) dy$  where  $C$  is the semi-ellipse  $\{x^2 + 4y^2 = 4, y \geq 0\}$  from the point  $(2,0)$  to the point  $(-2,0)$ .
- 7) Answer the following:
- Prove that the area bounded by a simple closed curve  $C$  is given by  $\frac{1}{2} \oint_C xdy - ydx$ .
  - Compute the area of the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1$ , using the formula  $A = \frac{1}{2} \oint_C xdy - ydx$ .

**Answer Key:**

- The common value is 0.5
- The common value is 0.8
- The common value is 8
- $1.5\pi$
- $0.5 \sin 64$
- $\frac{8}{3} + 2\left(e^4 - \frac{1}{e^4}\right)$
- a. Refer to the video    b.  $\pi ab$